

## RESEARCH ARTICLE

The Effects of Infant Births on Male–Female Relationships in *Cebus capucinus*

CLAIRE R. SHELLER\*, ZDANNA KING, AND KATHARINE JACK  
 Department of Anthropology, Tulane University, New Orleans, Louisiana

Most primates are characterized by cohesive male–female bonds that are maintained year round. While recent studies have addressed the selective pressures influencing the evolution of male–female relationships in primates, we know relatively little about the proximate mechanisms affecting them. It has been demonstrated that newborn white-faced capuchins (*Cebus capucinus*) attract the attention of other group members and this may be an important mechanism influencing male–female relationships. We studied two groups of *C. capucinus* in the Area de Conservación Guanacaste, Costa Rica, between February and July 2007. A total of 348 hr of focal data were collected on all adult males ( $n = 6$ ) residing in each of the study groups. During our study, 13 of the 14 group females were either pregnant or lactating, and 9 infants were born. We calculated an average daily affiliation rate between all group males combined and each adult female four weeks before and four weeks after the birth of her infant. Repeated measures ANOVAs revealed no significant changes in affiliation following infant births ( $F = 2.262$ ,  $df = 1$ ,  $P < 0.176$ ). Results remained nonsignificant for rank ( $F = 1.550$ ,  $df = 1$ ,  $P < 0.260$ ) and group membership ( $F = 0.729$ ,  $df = 1$ ,  $P < 0.429$ ). Infant sex was the only variable with a significant effect on affiliation rates between males and females ( $F = 10.020$ ,  $df = 1$ ,  $P < 0.019$ ). Adult males increased their affiliation with all adult females that gave birth to male infants ( $n = 4$ ), while their rates decreased with all but one of the adult females with female infants ( $n = 4$ ). While preliminary, these results indicate that the adult males may cultivate relationships with other males at a young age. *Am. J. Primatol.* 71:380–383, 2009. © 2009 Wiley-Liss, Inc.

**Key words:** affiliation; parturition; socialization; dispersal

## INTRODUCTION

A number of researchers have characterized the patterns of male–female social relationships in *Cebus* [Fedigan, 1993; O'Brien, 1991; Rose and Fedigan, 1995], but we are aware of no study that has explicitly examined how infant births affect these relationships. Newborn white-faced capuchins (*Cebus capucinus*) garner considerable attention from other group members [Manson, 1999]. Adult males take particular interest in new infants, often following the mother in order to groom, nuzzle or carry the new baby [MacKinnon, 2002]. This attraction to the newborn should also affect the male's relationship with new mothers, as the baby serves as a focal point to bring the sexes together. In this article, we examine how rates of affiliation among males and females change with the birth of infants.

The attraction of group males to new infants in primates may function to increase infant survivorship by providing, either directly or indirectly, protection from predators [Janson, 1986], increased access to food [Escobar-Páramo, 1989], protection from intragroup aggression [Escobar-Páramo, 1989; Ferreira et al., 2006; Hector et al., 1989; Kleindorfer and Wasser, 2004; Vessey and Meikle, 1984] and protection from infanticide [Busse and Gordon, 1984;

Vogel and Fuentes-Jiménez, 2006]. In some species, adult males also play an important role in infant socialization by encouraging appropriate behaviors, discouraging inappropriate behaviors, and by assisting in the forging of social bonds within the group [Burton, 1972; MacKinnon, 2002; Stein and Stacey, 1981].

The attention that adult males give to infants and their mothers may be dependent on a number of factors. Several studies have noted that, in some species, males are more attracted to male infants than female infants [Ogawa, 1995; Paul et al., 1996; Zhao, 1996]. Female rank may also play a role in the attractiveness of infants. Although adult females typically interact with infants of lower ranking females [O'Brien, 1991; Silk, 1999], in some species

Contract grant sponsor: Tulane's Committee on Research and Research Enhancement Fund.

\*Correspondence to: Claire R. Sheller, Department of Anthropology, Tulane University, 7041 Freret Street, New Orleans, LA 70118. E-mail: clairesheller@gmail.com

Received 26 September 2008; revised 11 December 2008; revision accepted 16 December 2008

DOI 10.1002/ajp.20661

Published online 23 February 2009 in Wiley InterScience (www.interscience.wiley.com).

it has been found that the adult males are more attracted to infants of high-ranking females (e.g. *Macaca sylvanus*: Paul et al., 1996; Small, 1990; and *Papio cynocephalus*: Stein, 1984]). Group composition can also affect patterns of affiliation simply because smaller groups have fewer available partners with whom individuals can form relationships. Males may also associate more with infants they have potentially sired or with females they have mated with in the past [Borries et al., 1999; Martin-Ordas and Colmenares, 2005; Menard et al., 2001].

In this study, we compare male–female affiliation levels for a 1-month period before and after the infant births in two groups of *C. capucinus*. We further examine how infant sex, female rank and group membership may influence changes in affiliation patterns. While we hope to examine the effect of infant paternity on these relationships in the future, genetic data are unavailable at this time.

## METHODS

### Study Site

We conducted our study in the Santa Rosa Sector of the Area de Conservación Guanacaste, Costa Rica. Santa Rosa encompasses approximately 108 km<sup>2</sup> of tropical dry forest at varying stages of regeneration. The park experiences a distinct wet season from late May to early December, with minimal rainfall during the remainder of the year [Fedigan and Jack, 2001].

We observed two groups of white-faced capuchins from February through July of 2007. Guanacaste (GC) was comprised of 21 members: 3 adult males, 9 adult females, 7 juveniles, 2 independent infants, and 5 infants that were born during the study. Los Valles (LV) was comprised of 15 members: 3 adult males, 5 adult females, 6 juveniles, 1 independent infant and 4 infants that were born over the course of the study. All infants were born during the dry season, between late March and early June 2007. All females who gave birth during the study period were pregnant when observations began. One LV infant died within days of its birth and its mother was excluded from analysis. At the time of this study, all males had maintained residence in their respective study groups for a minimum of one year.

### Data Analysis

Groups were observed for the majority of daylight hours spanning from 6:00 to 18:00, two to three times per week. A total of 348 hr of focal animal data were collected on adult males ( $\geq 10$  years) ( $n = 6$ ) during 10 min focal animal follows [Altmann, 1974]. This study complied with protocols approved by Tulane University's IACUC and the legal requirements of Costa Rica.

All durational, affiliative behaviors (including grooming, contact behaviors, and proximity behaviors within 2 m) were combined into a single affiliation category. Only those females who gave birth during the study period were included in the analysis. For each male–female dyad in the two study groups ( $n = 23$ ), we calculated an average daily affiliation rate for 4 weeks preceding (the “before” category) and 4 weeks following (the “after” category) the infant's birth. We also combined the average daily affiliation rates of all males residing in each group for each group female and examined how these changed after infant births. It should be noted that one male–female dyad (BB-MZ) from the LV group was not considered in our analyses. Data collection on MZ did not begin until he was elevated to full adult status on March 19. The female BB gave birth to her infant before March 19 and as such there is no “before” data available. This second data set was examined using a repeated measures ANOVA in SPSS to test whether the birth of an infant lead to any significant changes in affiliation patterns. We used the same test to examine the influence of female rank on affiliation patterns after the birth of an infant. In this case, we compared the affiliation rates in the alpha-female/all-male category (see Table I) with the rates from the other females/all-male category within the same group. Females were assigned a rank of either alpha or beta based on ad libitum observation of female feeding priority and instances of agonism and submission within each group. Our rank assignments were supported by concurrently collected data on females by other researchers (Mackenzie Bergstrom, personal communication). Next, we used a repeated measures ANOVA to determine if group membership had an effect on male–female affiliation rates after the infant births. We did this by comparing the mean affiliation rates of the female/all-male category in GC with those of LV, before and after infant births. Finally, we used the same test to determine if infant sex affected the male–female affiliation rates. This was done by comparing means of affiliation rates for the female/all-male categories of females with male infants with those of females with female infants, before and after births.

## RESULTS

Dyadic affiliation rates ranged from a low of 0.1% to a high of 11% during the before period and 0.4% and 14.1% during the after period. Fourteen of 23 male–female dyads displayed an increase in daily average affiliation, eight displayed a decrease and one displayed no change after the birth of an infant (Table I). Combining all group male affiliation data with each group female showed that the affiliation rates increased during the “after” period for five of the eight females, while they decreased for three of the females. The change in affiliation patterns before

**TABLE I. Comparative Affiliation Rates (%) for Male-Female Dyads ( $n = 23$ ) Before and After Infant Births**

Group	Dyad	Before	After	Change
GC	FL-BG*	0.1	3.2	+3.1
GC	FL-AD	1.1	0.9	-0.3
GC	FL-MM	3.1	8.3	+5.2
GC	FL-ALL MALES	1.5	3.8	+2.3
GC	LL-BG*	1.8	5.4	+3.6
GC	LL-AD	3.3	4.6	+1.3
GC	LL-MM	1.9	6.7	+4.7
GC	LL-ALL MALES	2.4	5.7	+3.3
GC	MV-BG*	3.3	4.1	+0.8
GC	MV-AD	2.7	13.8	+11.1
GC	MV-MM	0.5	6.0	+5.4
GC	MV-ALL MALES	2.3	8.1	+5.8
GC	**MX-BG*	9.3	14.1	+4.8
GC	**MX-AD	10.2	4.3	-5.9
GC	**MX-MM	2.7	1.0	-1.7
GC	**MX-ALL MALES	7.0	6.9	-0.2
GC	RM-BG*	7.2	4.4	-2.9
GC	RM-AD	0.7	7.9	+7.2
GC	RM-MM	1.8	3.8	+2.1
GC	RM-ALL MALES	3.1	5.2	+2.1
LV	BB-CY*	1.2	6.0	+4.9
LV	BB-WW	0.9	9.0	+8.1
LV	BB-ALL MALES	1.1	7.5	+6.4
LV	**KL-CY*	8.1	3.7	-4.4
LV	**KL-WW	3.4	3.0	-0.4
LV	**KL-MZ	3.9	5.1	+1.3
LV	**KL-ALL MALES	5.3	4.1	-1.2
LV	DL-CY*	11.0	4.6	-6.4
LV	DL-WW	5.4	0.4	-5.0
LV	DL-MZ	3.0	3.0	0.0
LV	DL-ALL MALES	6.5	2.7	-3.8

Alpha male indicated by “\*”, alpha female by “\*\*”, an increase in affiliation by “+”, and a decrease in affiliation by “-”.

and after the birth of an infant was nonsignificant ( $F = 2.262$ ,  $df = 1$ ,  $P < 0.176$ ). Next, we examined if group composition, female rank and infant sex had any effect on the changes in affiliation patterns after the infant births. Results remained nonsignificant with respect to group composition ( $F = 0.729$ ,  $df = 1$ ,  $P < 0.429$ ) and female rank ( $F = 1.550$ ,  $df = 1$ ,  $P < 0.260$ ), but infant sex had a significant effect on the affiliation rates between males and females ( $F = 10.020$ ,  $df = 1$ ,  $P < 0.019$ ). Group males increased their affiliation with females that gave birth to male infants ( $n = 4$ ), and decreased their affiliation with three out of four females that gave birth to female infants (Fig. 1). Each of the females who gave birth to the first infant in their group, both of which were male infants, experienced the most pronounced increase in affiliation rates with group males.

## DISCUSSION

Our study did not support the idea that, overall, the birth of an infant significantly alters male and




Fig. 1. Changes in affiliation rates (%) between adult females and all adult males after the birth of an infant in each group. Alpha female is indicated by “\*\*”, females with male infants are denoted by black bars, and females with female infants are shown by gray bars. Females are listed left to right in the order in which they gave birth. For example, in GC group, MV was the first to give birth, then FL, then MX, and so on.

female affiliative interactions. Roughly, half of the male–female dyads displayed an increase in affiliation after the birth of an infant while the other half displayed a decrease. Neither the rank of the mother nor group membership significantly influenced the attention that males gave to a female and her new infant. However, our finding that adult males were significantly more interested in females with male infants supports the findings from other species [*M. sylvanus*: Paul et al., 1996; *M. thibetana*: Ogawa, 1995; Zhao, 1996]. It is possible that birth order may explain this increase in affiliation because the first infants born into each study group were male and their mothers experienced the greatest increase in their affiliation rates. However, upon examination of the pattern of change in affiliation (see Fig. 1) for our largest study group (GC), birth order does not appear to be playing a role. It will be necessary to explore this possible effect with a larger data set. While our results are still preliminary, it is possible that the interest adult males have in male infants serves as an important start in their socialization process. In white-faced capuchins, the formation of cooperative bonds among males is important for group defense and establishing and maintaining group membership [Perry, 1998; Jack, 2003]. Jack and Fedigan [2004a,b] suggest that in this species, the formation and maintenance of cooperative bonds among group males are possible because of the high frequency of parallel dispersal (dispersing with group mates or into groups containing familiar individuals). Male white-faced capuchins emigrate from their natal group at four years of age on average, but have been documented to disperse as early as 20 months. This primary dispersal is often triggered by

the departure of adult males from the group, i.e. young males will follow older males when they disperse [Jack and Fedigan, 2004a]. Given that the average tenure length for adult males is four years [Jack and Fedigan, 2004b], it is possible that the infant to whom an adult male has bonded may indeed become a future dispersal partner. These males may either disperse together or the maturing natal male may later immigrate into the group containing the familiar older male. The results presented here provide the first tantalizing evidence that the formation of bonds between males and potential future dispersal partners may begin at a very young age.

## ACKNOWLEDGMENTS

This study was funded by a grant awarded to Dr. Katharine Jack through Tulane University's Research Enhancement Fund. Thanks to Amanda Melin, Andrew Childers, Fernando Campos, Laura Weckman, MacKenzie Bergstrom, Nigel Parr, Teresa Holmes, and Valerie Schoof for help with data collection and valuable input, and to Dr. David Corey and Margo Sidell for assistance with statistical analysis. Preparation of this manuscript was supported by a summer fellowship from Tulane's Committee on Research. Our research complied with protocols approved by Tulane University's IACUC and Costa Rican laws.

## REFERENCES

- Altmann J. 1974. Observational study of behavior: sampling methods. *Behavior* 49:227–265.
- Borries C, Launhardt K, Epplen C, Epplen JT, Winkler P. 1999. Males as infant protectors in Hanuman langurs (*Presbytis entellus*) living in multimale groups—defense pattern, paternity and sexual behaviour. *Behav Ecol Sociobiol* 46:350–356.
- Burton FD. 1972. The integration of biology and behavior in the socialization of *Macaca sylvanus* of Gibraltar. In: Poirer MA, editor. *Primate socialization*. New York: Random House. p 29–62.
- Busse CD, Gordon TP. 1984. Infant carrying by adult male mangabeys (*Cercocebus atys*). *Am J Primatol* 6:133–141.
- Escobar-Páramo P. 1989. The development of the wild black-capped capuchins (*Cebus apella*) in La Macarena, Colombia. *Field Studies of New World Monkeys, La Macarena, Colombia* 2:45–56.
- Fedigan LM. 1993. Sex differences and intersexual relations in adult white-faced capuchins (*Cebus capucinus*). *Int J Primatol* 14:853–877.
- Fedigan LM, Jack KJ. 2001. Neotropical primates in a regenerating Costa Rican dry forest: a comparison of howler and capuchin population patterns. *Int J Primatol* 22: 689–713.
- Ferreira RG, Izar RG, Lee PC. 2006. Exchange, affiliation, and protective interventions in semifree-ranging brown capuchin monkeys (*Cebus apella*). *Am J Primatol* 68:765–776.
- Hector AC, Seyfarth RM, Raleigh MJ. 1989. Male parental care, female choice and the effect of an audience in vervet monkeys. *Anim Behav* 38:262–271.
- Jack KM. 2003. Explaining variation in affiliative relationships among male white-faced capuchins (*Cebus capucinus*). *Folia Primatol* 74:1–16.
- Jack KM, Fedigan LM. 2004a. Male dispersal patterns in white-faced capuchins, *Cebus capucinus*, part 1: patterns and causes of natal emigration. *Anim Behav* 67:761–769.
- Jack KM, Fedigan LM. 2004b. Male dispersal patterns in white-faced capuchins, *Cebus capucinus*, part 2: patterns and causes of secondary dispersal. *Anim Behav* 67:771–782.
- Janson CH. 1986. Capuchin counterpoint. *Nat Hist* 95:44–53.
- Kleindorfer S, Wasser SK. 2004. Infant handling and mortality in yellow baboons (*Papio cynocephalus*) evidence for female reproductive competition. *Behav Ecol Sociobiol* 56:328–337.
- MacKinnon KC. 2002. Social development of wild white-faced capuchin monkeys (*Cebus capucinus*) in Costa Rica: an examination of social interactions between immatures and adult males [dissertation]. Berkeley (CA): University of California. Available from main library at University of California Berkeley; 308t 2002 341.
- Manson JH. 1999. Infant handling in wild *Cebus capucinus*: testing bonds between females? *Anim Behav* 57:911–921.
- Martin-Ordas G, Colmenares F. 2005. Social relationships among adult males and immature individuals in a mixed colony of baboons (*Papio* spp.): paternity certainty, sexual selection or female choice [abstract]? *Folia Primatol* 76:57–58.
- Menard N, von Segesser F, Scheffrahn W, Pastorini J, Vallet D, Gaci B, Martin RD, Gautier-Hion A. 2001. Is male-infant caretaking related to paternity and/or mating activities in wild Barbary macaques (*Macaca sylvanus*). *C R Acad Sci III* 324:601–610.
- O'Brien TG. 1991. Female–male social interactions in wedge-capped capuchin monkeys: benefits and costs of group living. *Anim Behav* 41:555–567.
- Ogawa H. 1995. Triadic male–female–infant relationships and bridging behaviour among Tibetan macaques (*Macaca thibetana*). *Folia Primatol* 64:153–157.
- Paul A, Kuester J, Arnemann J. 1996. The sociobiology of male–infant interactions in barbary macaques (*Macaca sylvanus*). *Anim Behav* 51:155–170.
- Perry S. 1998. Male–male social relationships in wild white-faced capuchins, *Cebus capucinus*. *Behavior* 135:139–172.
- Rose LM, Fedigan LM. 1995. Vigilance in white-faced capuchins, *Cebus capucinus*, in Costa Rica. *Anim Behav* 49:63–70.
- Silk JB. 1999. Why are infants so attractive to others? The form and function of infant handling in bonnet macaques. *Anim Behav* 57:1021–1032.
- Small MF. 1990. Alloparental behaviour in Barbary macaques, *Macaca sylvanus*. *Anim Behav* 39:297–306.
- Stein DM. 1984. *The sociobiology of infant and adult male baboons*. Norwood (NJ): Ablex Publishing Corporation. 229p.
- Stein DM, Stacey PB. 1981. A comparison of infant–adult male relations in a one-male group with those in a multi-male group for yellow baboons (*Papio cynocephalus*). *Folia Primatol* 36:264–276.
- Vessey SH, Meikle DB. 1984. Free-living rhesus monkeys: adult male interactions with infants and juveniles. In: Taub MT, editor. *Primate paternalism*. New York: Van Nostrand Reinhold Company Inc. p 113–126.
- Vogel ER, Fuentes-Jiménez A. 2006. Rescue behavior in white-faced capuchin monkeys during an intergroup attack: support for the infanticide avoidance hypothesis. *Am J Primatol* 68:1012–1016.
- Zhao QK. 1996. Male–infant–male interactions in Tibetan macaques. *Primates* 37:135–143.